

Specimen Label



Snapshot* 2.5 TG

Specialty Herbicide

*Trademark of Dow AgroSciences LLC

A selective preemergence herbicide for control of certain broadleaf weeds and annual grasses in:

- Landscape Ornamentals
- Christmas Tree Plantations
- Container Grown Ornamentals
- Field Grown Ornamentals
- Ground Covers / Perennials
- Non-bearing Fruit and Nut Trees
- Non-bearing Vineyards
- Non-cropland

Active Ingredients:

trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine	2.0%
isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5- isoxazolyl]-2,6-dimethoxybenzamide and isomers.....	0.5%
Inert Ingredients	97.5%
Total	100.0%

Contains 1.25 pounds active ingredient per 50 pound bag.

U.S. Patents 4,636,243 and 5,086,184.

EPA Reg. No. 62719-175

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Harmful If Swallowed Or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reaction In Some Individuals

Avoid breathing dust or spray mist and contact with eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

If swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for further treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements of this label about personal protective equipment (PPE), and restricted entry interval. The requirements in the box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For applications in landscape settings and in non-cropland sites, do not enter or allow others to enter the treated area until dusts have settled.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container only. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, in accordance with applicable regulations, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Instructions and Information

Snapshot* 2.5 TG herbicide is a preemergence product for control of certain broadleaf weeds and annual grasses in container and landscape ornamentals, nursery stock, Christmas trees, ground covers/perennials, non-bearing fruit and nut trees, non-bearing vineyards and non-cropland areas. Apply Snapshot 2.5 TG anytime prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary with rate of Snapshot 2.5 TG applied, weed population, potting media or soil conditions, temperature, watering regime, and other factors. Following application, user should monitor and observe level of weed control over time to determine when additional applications may be needed.

General Use Precautions and Restrictions

In Arizona: The state of Arizona has not approved Snapshot* 2.5 TG specialty herbicide for use on plants grown for agricultural/commercial production; such as on Christmas tree plantations, container grown or field grown ornamentals, non-bearing fruit and nut trees and non-bearing vineyards.

Snapshot 2.5 TG controls weeds growing from seed. Snapshot 2.5 TG does not control established weeds, weeds growing from stolens, rhizomes, or root pieces. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation is required to activate Snapshot 2.5 TG. Optimum weed control is obtained when Snapshot 2.5 TG is activated within 3 days of application. If rainfall or irrigation does not occur within 3 days of application and tillage is possible, Snapshot 2.5 TG may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1-2 inches of soil. Failure to activate Snapshot 2.5 TG within 3 days of application may result in erratic control of annual grasses. Do not apply when wind conditions favor drift of Snapshot 2.5 TG granules from the target area.

Repeat applications at 150 pounds per acre and higher should not be made sooner than 60 days after a previous application of Snapshot 2.5 TG. Do not apply over 600 pounds per acre total of Snapshot 2.5 TG within a 12 month period.

Tolerance of Turfgrass Adjacent to Ornamental Plantings: Accidental application to turf may occur when applying Snapshot 2.5 TG to ornamental plantings. Snapshot 2.5TG is not recommended for weed control in turfgrasses, but turfgrasses are generally tolerant to small amounts of this product that fall outside of the intended area of application.

Do not aerially apply Snapshot 2.5TG.

Treatment Species Not Listed on the Label for Snapshot 2.5 TG

Users who wish to use Snapshot 2.5 TG on plant species not recommended on this label may determine the suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury for during 30 to 60 days of normal growing conditions to determine if the treatment is safe to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of Snapshot 2.5 TG on plant species not recommended on this label.

Application Instructions

Apply Snapshot 2.5 TG using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or crop injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

Application Techniques for Applying Snapshot 2.5 TG

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.
- The weight-to-volume conversion table below provides equivalent amounts of Snapshot 2.5 TG.

Snapshot 2.5 TG Broadcast Rates

Rate lb/Acre	Amount/1000 sq ft		Amount/100 sq ft	
	(lb)	(quarts)	(lb)	(cups)
100	2.30	1.75	0.23	0.7
150	3.45	2.60	0.35	1.0
200	4.60	3.50	0.46	1.4

One pound of Snapshot 2.5 TG = 0.75 quart = 3 cups

Spreader Settings as a Guide for Calibration

Note: These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by Dow AgroSciences or the spreader manufacturer.

Speed (mph)	Settings for Warren T-7 II Spreader		
	Target Rate Of Snapshot 2.5 TG		
	100 lb/acre	150 lb/acre	200 lb/acre
2.0	2-1/2	3-1/4	3-3/4
2.5	3	3-1/2	4-1/4
3.0	3-1/4	4	5

Warren spreader settings are displayed in half-number increments only. The suggested settings that indicate quarter settings can be accomplished by placing the calibration arm between the half number marks on the spreader.

Weeds Controlled or Suppressed by Snapshot 2.5 TG

Weeds Controlled When Applied at 100 lb/acre (2.3 lb/1000 sq ft)

Common Name	Scientific Name
aster, slender	<i>Aster exilis</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
bluegrass, annual	<i>Poa annua</i>
bursage, annual	<i>Ambrosia acanthicarpa</i>
celery, wild	<i>Apium leptophyllum</i>
chickweed, common	<i>Stellaria media</i>
clover white	<i>Trifolium repens</i>
crabgrass	<i>Digitaria</i> spp.
cudweed, purple	<i>Gnaphalium purpureum</i>
cupgrass, southwestern	<i>Eriochloa gracilis</i>
fiddleneck, coast	<i>Amsinckia intermedia</i>
filaree, redstem	<i>Erodium cicutarium</i>
fleabane, blackleaved	<i>Conyza bonariensis</i>
fleabane, dwarf	<i>Conyza ramosissima</i>
foxtail, yellow	<i>Setaria glauca</i>
groundcherry, lanceleaf	<i>Physalis lanceifolia</i>
henbit	<i>Lamium amplexicaule</i>
horseweed	<i>Conyza canadensis</i>
junglerice	<i>Echinochloa colonum</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambsquarters, common	<i>Chenopodium album</i>
mallow, little	<i>Malva parviflora</i>
mustard, Indian	<i>Brassica juncea</i>
mustard, wild	<i>Sinapis arvensis</i>
nightshade, black	<i>Solanum nigrum</i>
oat, wild	<i>Avena fatua</i>
panicum, fall	<i>Panicum</i> <i>dichotomiflorum</i>
pepperweed, Virginia	<i>Lepidium virginicum</i>
pigweed	<i>Amaranthus</i> spp.
pineappleweed	<i>Matricaria</i> <i>matricarioides</i>
plantain, slender	<i>Plantago elongata</i>
purslane, common	<i>Portulaca oleracea</i>
radish, wild	<i>Raphanus raphanistrum</i>
ragweed, common	<i>Ambrosia artemisiifolia</i>
rocket, London	<i>Sisymbrium irio</i>
rockpurslane, desert	<i>Calandrinia ciliata</i>
shepherdspurse	<i>Capsella bursa-pastoris</i>
sibara	<i>Sibara virginica</i>
smartweed, Pennsylvania	<i>Polygonum</i> <i>pensylvanicum</i>
sowthistle, annual	<i>Sonchus oleraceus</i>
speedwell, purslane	<i>Veronica peregrina</i>
sprangletop, Mexican	<i>Leptochloa univervia</i>
telegraphplant	<i>Heterotheca grandiflora</i>
thistle, Russian	<i>Salsola iberica</i>
witchgrass	<i>Panicum capillare</i>

In Addition to the Weeds Listed at the 100 lb/acre Rate, the Following Weeds Will Be Controlled at 150 lb/acre (3.5 lb/1000 sq ft)

Common Name	Scientific Name
aster, heath	<i>Aster ericoides</i>
bittercress	<i>Cardamine oligosperma</i>
bittercress, hairy	<i>Cardamine hirsuta</i>
brassbuttons, southern	<i>Cotula australis</i>
carrot, wild	<i>Daucus carota</i>
chamberbitter	<i>Phyllanthus urinaria</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
dandelion	<i>Taraxacum officinale</i>
eclipta	<i>Eclipta prostrata</i>
fireweed	<i>Erechtites hieracifolia</i>
foxtail, giant	<i>Setaria faberi</i>
galinsoga, hairy	<i>Galinsoga ciliata</i>
geranium, Carolina	<i>Geranium carolinianum</i>
groundsel, common	<i>Senecio vulgaris</i>
ladysthumb	<i>Polygonum persicaria</i>
lettuce, prickly	<i>Lactuca serriola</i>
lovegrass	<i>Eragrostis</i> spp.
mallow, dwarf	<i>Malva rotundifolia</i>
maretail	<i>Hippuris vulgaris</i>
mayweed	<i>Anthemis cotula</i>
morningglory, ivyleaf	<i>Ipomoea hederacea</i>
mustard, black	<i>Brassica nigra</i>
pennywort	<i>Hydrocotyle</i> spp.
phyllanthus, long-stalk	<i>Phyllanthus tenellus</i>
plantain, bracted	<i>Plantago aristata</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
pokeweed, common	<i>Phytolacca americana</i>
rockpurslane, redmaids	<i>Calandrinia ciliata</i> var. <i>menziesii</i>
ryegrass, annual	<i>Lolium multiflorum</i>
sida, prickly	<i>Sida spinosa</i>
sorrell, red	<i>Rumex acetosella</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, hyssop	<i>Euphorbia hyssopifolia</i>
spurge, spotted	<i>Euphorbia maculata</i>
sweetclover, yellow	<i>Melilotus officinalis</i>
tansymustard, green	<i>Descurainia pinnata</i>
velvetleaf	<i>Abutilon theophrasti</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

In Addition to the Weeds Listed at the 100 lb/acre and 150 lb/acre Rates, the Following Weeds Will Be Controlled at 200 lb/acre (4.6 lb/1000 sq ft)

Common Name	Scientific Name
barley, hare	<i>Hordeum leporinum</i>
bromegrass	<i>Bromus</i> spp.
burclover, California	<i>Medicago polymorpha</i>
cheat	<i>Bromus secalinus</i>
datura	<i>Datura</i> spp.
dogfennel	<i>Eupatorium capillifolium</i>
eveningprimrose	<i>Oenothera</i> spp.
fescue, rattail	<i>Vulpia myuros</i>

filaree, whitestem
goosefoot, nettleleaf
goosegrass
jimsonweed
knotweed, silversheath

kochia
medic, black
mullein, turkey
nettle, burning
nettle, stinging
ox tongue, bristly
pimpernel, scarlet
sandbur, field
signalgrass
sowthistle, spiny
spurge, petty
spurge, prostrate
stinkgrass
sunflower
swinecress
thistle, musk
willoweed, panicle
woodsorrel, creeping

Erodium moschatum
Chenopodium murale
Eleusine indica
Datura stramonium
Polygonum argyrocoleon
Kochia scoparia
Medicago lupulina
Eremocarpus setigerus
Urtica urens
Urtica dioica
Picris echioides
Anagallis arvensis
Cenchrus incertus
Brachiaria spp.
Sonchus asper
Euphorbia pepplus
Euphorbia humistrata
Eragrostis cilianensis
Helianthus spp.
Coronopus didymus
Carduus nutans
Epilobium paniculatum
Oxalis corniculata

In Addition to the Weeds Controlled, the Following Weeds Will Be Partially Controlled or Suppressed at 200 lb/acre (4.6 lb/1000 sq ft)

Common Name	Scientific Name
bindweed, field	<i>Convolvulus arvensis</i>
carpetweed	<i>Mollugo verticillata</i>
dock, curly	<i>Rumex crispus</i>
johnsongrass (seedling)	<i>Sorghum halepense</i>
mallow, Venice	<i>Hibiscus trionum</i>
milkweed, honeyvine	<i>Ampelamus albidus</i>
morningglory, tall	<i>Ipomoea purpurea</i>
panicum, Texas	<i>Panicum texanum</i>
pusley, Florida	<i>Richardia scabra</i>
shattercane	<i>Sorghum bicolor</i>

Labeled Uses

Ornamental Plantings

Snapshot 2.5 TG is recommended as a preemergence treatment for control of certain broadleaf weeds and annual grasses in container grown ornamentals, landscape ornamentals, field grown ornamentals, ground covers/perennials, Christmas trees, non-bearing fruit and nut crops, and non-bearing vineyards. Apply Snapshot 2.5 TG anytime prior to germination of target weeds, or immediately after cultivation.

Refer to the "General Information" section of this label for General Use Precautions and information on application, application rates and weeds controlled.

Special Use Precautions

To avoid possible injury, do not apply Snapshot 2.5 TG to:

- nursery, forest, or Christmas tree: seedling beds, cutting beds, or transplant beds.
- nursery seedbeds or forest or Christmas tree seedling transplant beds.
- unrooted liners or cuttings that have been planted in pots for the first time.
- pots less than four inches wide.
- bedding plants or areas where bedding plants will be planted or transplanted within one year following application.
- ground covers until they are established and well rooted.

Do not apply Snapshot 2.5 TG to newly transplanted ornamentals, nursery stock, Christmas trees, ground covers, non-bearing fruit and nut trees and non-bearing vineyards until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury may occur.

Repeat applications at 150 pounds per acre and higher should not be made sooner than 60 days after a previous application of Snapshot 2.5 TG. Do not apply over 600 pounds per acre total of Snapshot 2.5 TG within a 12 month period.

Where loss of stand has occurred in field grown ornamentals, use untreated soil as fill around roots when replacing plants or injury may occur.

Note: Injury on the following plant species has been observed following applications of Snapshot 2.5 TG and use is not recommended.

Acanthus mollis (oak leaf acanthus)
Ajuga spp. (bugle)
Arenaria montana (mountain sandwort)
Brassica spp. (mustard)
Dicentra luxuriant (*luxuriant bleeding heart*)
Digitalis purpurea (foxglove)
Echinacea purpurea (purple coneflower)
Euonymus alatus 'compacta' (dwarf burning bush)
Euphorbia spp. (spurge)
Geranium cinereum 'Ballerina' (*ballerina cranesbill*)
Gypsophila paniculata (*white festival baby's breath*)
Hydrangea spp. (*hydrangea*)
Iberis spp. (*candytuft*)
Juniperus horizontalis 'Prince of Wales' (*Prince of Wales juniper*)
Melaleuca quinquenervia (*cajeput tree*)
Phlox paniculata (*summer phlox*)
Pulmonaria longifolia (*Bertram Anderson lingwort*)
Rhododendron carolinianum (*Carolina rhododendron*)
Rhododendron catawbiense 'Roseum elegans' (*Roseum elegans rhododendron*)
Sedum spp. (*stonecrop*)
Vinca minor 'Atropurpurea' (*wine periwinkle*)
Yucca recurvifolia (*green yucca*)

Snapshot 2.5 TG May Be used On the Following Established Plant Species

(Note: Limitations on recommended treatment methods)

Trees

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Abies balsamea</i>	Fir, balsam	C, F
<i>Abies concolor</i>	Fir, white	F
<i>Abutilon hybridum</i>	Albus-flowering maple	C, F
	Luteus-flowering maple	C, F
	Roseus-flowering maple	C, F
	Tangerine-flowering maple	C, F
	Vesuvius red-flowering maple	F
<i>Acer ginnmala</i>	Flame maple	F
<i>Acer rubrum</i>	Red maple	F
	Red sunset maple	F
<i>Acer saccharinum</i>	Silver maple	C, F
<i>Alsophila australis</i>	Australian tree fern	C, F
<i>Areacastrum romanzoffianum</i>	Queen palm	C, F
<i>Betula nigra</i>	Birch, river	C, F
<i>Betula papyrifera</i>	Paper birch	F
<i>Brachychiton populneus</i>	Bottle tree	C, F
<i>Bucida buceras</i>	Black olive	F
<i>Ceratonia siliqua</i>	Carob	F
<i>Cercis canadensis</i>	Redbud	C, F
<i>Chamaecyparis obtusa</i>	Filicoides-fernspray cypress	F
<i>Chamaecyparis obtusa</i>	Gracilis-slender Hinoki cypress	F
<i>Chamaecyparis pisifera</i>	Sawara-false cypress	F
	Squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	Cat Palm	F
<i>Chamaedorea costaricana</i>	Palm	C, F
<i>Chamaedorea elegans</i>	Palm	C, F
<i>Cornus florida</i>	Parlor palm	C, F
	Cloud nine dogwood	C, F
	Dogwood, flowering	C, F
<i>Crataegus viridis</i>	Green hawthorn	F
<i>Cryptomeria japonica</i>	Cryptomeria, Japanese	C, F
<i>Cupaniopsis anacardioides</i>	Carrot wood	F
<i>Cupressocyparis x 'Emerald Isle'</i>	Emerald island leyland cypress	C, F
<i>Cupressus arizonica</i>	Arizona cypress	F
<i>Cupressus glabra</i>	Arizona cypress	F
<i>Cupressus sempervirens</i>	Italian cypress	C, F
<i>Dicksonia antarctica</i>	Tasmanian tree fern	C, F
<i>Elaeagnus angustifolia</i>	Russian olive	C, F
<i>Elaeagnus x 'Gilt edge'</i>	Gilt edge strawberry	C, F
<i>Eucalyptus camaldulensis</i>	Red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	Eucalyptus, mealy	F
	Silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i>	Coolibah tree	C, F
<i>Eucalyptus sideroxylon</i>	Eucalyptus, red ironbark	F
<i>Ficus benjamina</i>	Ficus	C, F
	Mini ficus	C, F
<i>Fraxinus udhei</i>	Shamel ash	C, F
<i>Ginkgo biloba</i>	Ginkgo (Maidenhair tree)	F

<i>Gleditsia triacanthos</i>	Honey locust	F
	Shademaster honey locust	F
<i>Heteromeles arbutiflora</i>	Toyon	F
<i>Illicium floridanum</i>	Florida anise-tree	C, F
<i>Juniperus virginiana</i>	Redcedar, Eastern	C, F
<i>Liquidambar styraciflua</i>	Sweetgum, American	F
<i>Magnolia grandiflora</i>	Magnolia, Southern	C, F
<i>Morus alba</i>	White mulberry	F
<i>Musa aluminata</i>	Banana	C, F
<i>Oxydendrum arboreum</i>	Sourwood	C, F
<i>Picea abies</i>	Pendula-weeping Norway spruce	C, F
	Repens-spreading Norway spruce	C, F
	Spruce, Norway	C, F
<i>Picea glauca</i>	Conica-dwarf Alberta spruce	C, F
<i>Picea glauca conica</i>	Dwarf Alberta spruce	F
<i>Picea pungens</i>	Dwarf globe blue spruce	C, F
	Glauca-Colorado blue spruce	C, F
	Hoopsii-Hoop's blue spruce	C, F
	Koster-Koster blue spruce	F
<i>Pinus aristata</i>	Bristlecone pine	F
<i>Pinus canariensis</i>	Canary Island pine	F
<i>Pinus contorta</i>	Shore pine, beach pine	F
<i>Pinus eldarica</i>	Eldarica pine	C, F
<i>Pinus leucodermis</i>	Bosnian pine	C, F
<i>Pinus mugo</i>	Pumilio-shrubby swiss mountain pine	C, F
<i>Pinus nigra</i>	Austrian black pine	C, F
<i>Pinus radiata</i>	Monterey pine	F
<i>Pinus strobus</i>	Eastern white pine	C, F
<i>Pinus strobus</i>	white pine	C, F
<i>Pinus sylvestris</i>	Columnar Scotch pine	C, F
	Scotch pine	C, F
<i>Pinus thunbergiana</i>	Japanese black pine	C, F
<i>Platanus occidentalis</i>	American sycamore	F
<i>Platanus racemosa</i>	Californina sycamore	F
<i>Podocarpus</i> spp.	Podocarpus	F
<i>Populus deltoides</i>	Cottonwood	F
<i>Prosopis chilensis</i>	Chilean mesquite	C, F
<i>Prunus yedoensis</i>	Yoshino flowering cherry	F
<i>Quercus ilicifolia</i>	Bear oak	F
<i>Quercus palustris</i>	Pin oak	F
<i>Quercus phellos</i>	Willow oak	C, F
<i>Quercus rubra</i>	Red oak	C, F
<i>Quercus virginiana</i>	Live oak	C, F
<i>Salix babylonica</i>	Babylon weeping willow	F
	Corkscrew willow	F
<i>Sequoiadendron giganteum</i>	Giant sequoia	F
<i>Swietenia mahogani</i>	Mahogany	F
<i>Tabebuia caraiba</i>	Yellow tab	F
<i>Tsuga canadensis</i>	Eastern hemlock	C, F
<i>Ulmus parvifolia</i>	Chinese elm	F
<i>Washingtonia robusta</i>	Mexican fan palm	F

Shrubs

Recommended Treatment Method

C = Container Grown

F = Field Grown

Scientific Name	Common Name	
<i>Abelia grandiflora</i>	Edward Goucher abelia	C, F
	Glossy abelia	C, F
<i>Acacia abyssinica</i>	Abyssinica acacia	C, F
<i>Acacia redolens</i>	Acacia, prostrate	C, F
<i>Acacia stenophylla</i>	Shoestring acacia	C, F
<i>Acalypha wilkesiana</i>	Copper leaf	C, F
<i>Acer ginnala</i>	Amur maple	C, F
<i>Acer palmatum</i>	Coral bark Japanese maple	C, F
	Dwarf Japanese maple	C, F
<i>Agapanthus africanus</i>	Queen Anne Lily of the Nile	C, F
<i>Agave americana</i>	Century plant	F
<i>Anisodonteia hypomandarum</i>	Cape mallow	C, F
<i>Aptenia cordifolia</i>	Red apple aptenia	C, F
<i>Ardisia japonica</i>	Chirimen marlberry	C, F
<i>Astilbe arendsii</i>	false spiraea	C, F
<i>Astilbe chinensis</i>	Astilbe/false spirea	C, F
<i>Athyrium nipponicum</i>	Japanese painted fern	C, F
<i>Baccharis pilularis</i>	coyotebush	F
<i>Berberis gladwynensis</i>	William Penn barberry	C, F
<i>Berberis mentorensis</i>	Mentor barberry	C, F
<i>Berberis thunbergii</i>	Aurea-golden Japanese barberry	F
	Crimson pygmy barberry	C, F
	Rose glow barberry	C, F
	Atropurea-redleaf Japanese barberry	F
	Cherry bomb barberry	C, F
<i>Bougainvillea</i> spp.	Barbara Karst	C, F
	California gold	C, F
	Pink pixie	C, F
	Scarlet O'Hara	C, F
	Temple fire	C, F
	Texas dawn	C, F
<i>Buxus x 'Green velvet'</i>	Green velvet boxwood	C, F
<i>Buxus microphylla japonica</i>	Boxwood, Japanese	C, F
<i>Buxus microphylla Koreana</i>	Korean boxwood	F
<i>Buxus sempervirens</i>	Boxwood, common	C, F
<i>Callistemon citrinus</i>	Bottlebrush, lemon	F
<i>Callistemon viminalis</i>	Weeping bottlebrush	C, F
<i>Calluna vulgaris</i>	spring torch scotch heather	C, F
<i>Camellia japonica</i>	Camellia	C, F
<i>Caryopteris x clandonen</i>	Dark knight bluebeard	C, F
<i>Cassia artemisioides</i>	Cassia, feathery	C, F
<i>Ceanothus</i> spp.	Wild lilac	F
<i>Cephalotaxus drupacae</i>	Plum yew	C, F
<i>Cerastium tomentosum</i>	Snow-in-summer	C, F
<i>Chamaecyparis obtusa</i>	Kosteri cypress	C, F
	Nana-dwarf Hinoki cypress	C, F
	Torulosa cypress	C, F
<i>Chamaecyparis pisifera</i>	baileyi-dogwood	F
	Flaviramea-dogwood	F
	Squarrosa Minima cypress	C, F
<i>Chamaecyparis pisifera</i> spp.	Filifera-thread cypress	C, F

<i>Chrysalidocarpus lutescens</i>	Areca palm	F	<i>Hibiscus syriacus</i>	Rose of Sharon,Red Bird	C, F
<i>Clethra alnifolia</i>	Summersweet	C, F		Rose of Sharon,Red Heart	F
<i>Cleyera japonica</i>	Cleyera, Japanese	C, F		Rose of Sharon,Woodbridge	C, F
<i>Coleonema pulchrum</i>	Pink breath of heaven	C, F		Rose of Sharon, Aphrodite	
<i>Convolvulus cneorum</i>	Bush morning glory	C, F	<i>Ilex aquifolium</i>	Rose of Sharon, Helene	
<i>Cornus alba</i>	Sibirica-Siberian dogwood	C, F		Balkans holly	F
<i>Cornus stolonifera</i>	Baileyi-red-osier dogwood	F	<i>Ilex aquipernyi</i>	Gold coast holly	F
	Flaviramea-yellowtwig dogwood	F	<i>Ilex attenuata</i>	San Jose holly	C, F
<i>Cotinus coggygria</i>	Royal purple smoke tree	C, F	<i>Ilex cornuta</i>	Savannah holly	C, F
<i>Cotinus dammeri</i>	Coral beauty smoke tree	C, F		Burford holly	C, F
	Eichholz smoke tree	C, F		dwarf Burford holly	C, F
<i>Cotoneaster adpressus</i>	Praecox-early cotoneaster	C, F	<i>Ilex crenata</i>	needle point holly	C, F
<i>Cotoneaster apiculatus</i>	Cotoneaster, cranberry	C, F		Compacta-dwarf Japanese holly	C, F
<i>Cotoneaster congestus</i>	Cotoneaster, Pyrenees	C, F	<i>Ilex crenata</i>	Convexa holly	C, F
<i>Cotoneaster dammeri</i>	Cotoneaster, bearberry	C, F		Dwarf Chinese holly	C, F
<i>Cotoneaster himalayana</i>	Himalayan cotoneaster	C, F		Green luster holly	C, F
<i>Cotoneaster horizontalis</i>	Cotoneaster, rock	C, F		Helleri-Heller's Japanese holly	C, F
<i>Cotoneaster opiculata</i>	Cotoneaster	C, F	<i>Ilex glabra</i>	Hetzii's Japanese holly	C, F
<i>Cycas revoluta</i>	Sago palm	C, F		Stokesii Japanese holly	C, F
<i>Cytisus praecox</i>	Hollandia-warminster broom	C, F	<i>Ilex meserveae</i>	compacta-compact inkberry holly	C, F
<i>Cytisus scoparius</i>	Lena-Scotch broom	C, F		Nordica-inkberry holly	C, F
<i>Cytisus spp.</i>	Holandia-Scotch broom	F		Blue boy holly	C, F
<i>Daphne odora</i>	Fragrant daphne	C, F		Blue girl holly	C, F
<i>Deutzia crenata</i>	Nakiana-dwarf deutzia	C, F		China boy holly	
<i>Deutzia gracilis</i>	Slender gracilis	C, F	<i>Ilex vomitoria</i>	China girl holly	
<i>Dodonea viscosa</i>	Hopseed bush	F		Ebony magic holly	F
<i>Elaeagnus pungens</i>	Fruitland silver berry	C, F		Nana-dwarf yaupon holly	C, F
<i>Erica cinerea</i>	Purple bell heather	C, F	<i>Illicium annisatum</i>	Pendula-weeping yaupon holly	C, F
<i>Erica vagans</i>	Cornish heather	C, F	<i>Itea ilicifolia</i>	yaupon holly	C, F
<i>Erica x darleyensa</i>	Mediterranean pink heather	C, F	<i>Ixora collinea</i>	Mystery gardenia	C, F
<i>Eugenia myrtifolia</i>	Dwarf brush cherry	C, F	<i>Juniperus chinensis</i>	Henry Garnet holly leaf sweetspire	C, F
<i>Euonymus x 'Aureo variegatus'</i>	Gold spot euonymus	C, F		Ixora	C, F
<i>Euonymus x 'Chollipo'</i>	Chollipo euonymus	C, F		hollywood juniper	C, F
<i>Euonymus fortunei</i>	Canadale gold euonymus	C, F		Media-old gold juniper	C, F
	Emerald'n gold euonymus	F		pfitzer juniper	C, F
	Sunspot euonymus	C, F		Pfizerana glauca-blue juniper	C, F
<i>Euonymus japonica</i>	Silver king euonymus	F	<i>Juniperus conferta</i>	Pfizerana-pfizer juniper	C, F
	Variegated evergreen euonymus	C, F		Sea green juniper	F
<i>Euonymus kiatschovica</i>	Spreading euonymus	C, F	<i>Juniperus horizontalis</i>	Torulosa-hollywood juniper	C, F
<i>Euonymus vegetus</i>	Bigleaf wintercreeper	C, F		Emerald sea shore juniper	C, F
<i>Euryops pectinatus</i>	Dwarf euryops	C, F		Shore juniper	C, F
<i>Fatsyhedera japonica</i>		C, F		Andorra juniper	C, F
<i>Fatsia japonica</i>	Japanese aralia	C, F		Bar Harbor juniper	C, F
<i>Felicia amelooides</i>	Blue marguerite	C, F		Blue chip juniper	C, F
<i>Forsythia intermedia</i>	Forsythia, border	C, F		Blue rug juniper	C, F
<i>Forsythia x 'Spring glory'</i>	Spring glory forsythia	C, F		Creeping juniper	C, F
<i>Gardenia jasminoides</i>	August beauty gardenia	C, F	<i>Juniperus procumbens</i>	Dwarf Andorra juniper	C, F
	Gardenia	C, F		Huntington blue juniper	C, F
	Radican gardenia	C, F		Plumosa-Andorra juniper	C, F
<i>Gaultheria shallon</i>	Salal/lemon leaf	C, F		Wiltonii-blue carpet juniper	C, F
<i>Gelsemium sempervirens</i>	Carolina jessamine	C, F	<i>Juniperus prostrata</i>	Nana-dwarf Japaneses garden juniper	C, F
<i>Genista pilosa</i>	Woadwaxen	C, F		Prostrata juniper	C, F
<i>Hibiscus rosa-sinensis</i>	Ross Estey-hibiscus	C, F			

<i>Juniperus sabina</i>	Broadmoor juniper	C, F	<i>Pinus mugo</i>	Mugo-mugho pine	C, F
	Foemina-Hicks juniper	C, F	<i>Pittosporum tobira</i>	Green pittosporum	C, F
	Savin juniper	C, F		Wheeler's dwarf pittosporum	C, F
	Tamariscifolia-Tam juniper	C, F	<i>Plumbago ariculata</i>	Blue cape plumbago	F
<i>Juniperus scopulorum</i>	Emerald green juniper	F	<i>Plumbago capensis</i>	Plumbago	C, F
<i>Juniperus squamata</i>	Blue juniper	C, F	<i>Podocarpus</i>	Yewpine	C, F
	Blue star juniper	C, F	<i>macrophyllus</i>		
	Parsonii juniper	C, F	<i>Polygala dalmaisiana</i>	Sweet pea shrub	C, F
<i>Kalmia latifolia</i>	Laurel, mountain	C, F	<i>Polystichum</i>	Tassel fern	C, F
<i>Lagerstroemia indica</i>	Crape myrtle	C, F	<i>polyblepharum</i>		
<i>Lantana</i> spp.	Lantana	C, F	<i>Potentilla fragiformis</i>	Cinquefoil	F
<i>Lavandula angustifolia</i>	English lavender	C, F	<i>Potentilla fruticosa</i>	Cinquefoil	C, F
<i>Lavandula latifolia</i>	English spike lavender	C, F		Gold drop potentilla	F
<i>Lavandula officianalis</i>	English lavender	C, F		Goldfinger potentilla	C, F
<i>Leptospermum scoparium</i>	New Zealand tea tree	C, F		Red ace potentilla	C, F
				Sunset potentilla	C, F
<i>Leucothoe axillaris</i>	Leucothoe, coast	C, F		Tangerine potentilla	C, F
<i>Leucothoe fontanesiana</i>	Leucothoe, drooping	C, F	<i>Potentilla verna</i>	Spring cinquefoil	C, F
<i>Ligustrum japonicum</i>	Privet, Japanese	C, F	<i>Prunus glandulosa</i>	Dwarf pink flowering almond	C, F
	wax ligustrum	C, F	<i>Pyracantha fortuneana</i>	Lolendei Monrovia pyracantha	C, F
	yellow tip ligustrum	C, F		Monon pyracantha	F
<i>Ligustrum lucidum</i>	Privet, glossy	C, F		Red elf hybrid pyracantha	F
<i>Ligustrum ovalifolium</i>	California privet	F		Rutgers hybrid pyracantha	C, F
<i>Ligustrum texanum</i>	Howardi privet	C, F		Santa Cruz pyracantha	C, F
	wax leaf privet	C, F		Victory pyracantha	F
<i>Ligustrum vicaryi</i>	Privet, golden	F	<i>Raphiolepis indica</i>	Charisma-Monruce raphiolepis	C, F
	Vicary golden privet	F		Enchantress-Moness raphiolepis	F
<i>Ligustrum vulgare</i>	Lodense privet	C, F		Raphiolepis (India hawthorn)	C, F
<i>Livistona chinensis</i>	Chinese fountain palm	F		Springtime-Monme raphiolepis	F
<i>Lonicera fragrantissima</i>	Winter honeysuckle	C, F	<i>Raphiolepis ovata</i>	Roundleaf raphiolepis	C, F
<i>Lonicera periclymenum</i>	Flowering woodbine	C, F	<i>Rhododendron</i>	Cannon's double azalea	C, F
	Serotina woodbine	C, F	<i>calendulaceum</i>	Flame azalea	F
<i>Lonicera sempervirens</i>	Trumpet honeysuckle	C, F		Golden flare azalea	C, F
<i>Loropetalum chinense</i>	Fringe flower	C, F		Klondike azalea	C, F
<i>Mahonia aquifolium compacta</i>	Dwarf Oregon grape	C, F	<i>Rhododendron campylocarpum</i>	Butterfly rhododendron	F
<i>Mahonia bealei</i>	Leather leaf mahonia	C, F	<i>Rhododendron carolinianum x daurium</i>		
<i>Mahonia repens</i>	Creeping mahonia	C, F	<i>Rhododendron catawbiense</i>	Catawba album rhododendron	C, F
<i>Myrica cerifera</i>	Wax myrtle	C, F		Catawba rhododendron	C, F
<i>Nandina domestica</i>	Compacta-dwarf heavenly bamboo	C, F		Lord Roberts rhododendron	C, F
	Harbour dwarf-heavenly bamboo	C, F		Rocket rhododendron	C, F
	Heavenly bamboo (Nandina)	C, F	<i>Rhododendron caucasicum x ponticum</i>	Cuninham White rhododendron	C, F
	Nana compacta-heavenly bamboo	C, F			
	Nana purpurea-heavenly bamboo	C, F	<i>Rhododendron exbury</i>	Cannon's double azalea	C, F
	Woods dwarf-heavenly bamboo	C, F		golden flare azalea	C, F
<i>Nerium oleander</i>	Hardy red oleander	C, F		Klondike azalea	C, F
	Oleander	C, F	<i>Rhododendron forrestii repens</i>	Gomer Waterer rhododendron	C, F
	Ruby lace oleander	C, F	<i>Rhododendron forrestii x griersonianum</i>		
<i>Osmanthus fortunei</i>	Fortunes osmanthus	C, F	<i>Rhododendron griffithianum</i>	Jean Marie rhododendron	C, F
<i>Pachysandra terminalis</i>	Japanese spurge	C, F			
<i>Phoenix roeloenii</i>	Pigmy date palm	C, F			
<i>Photinia fraseri</i>	Fraser's photinia	C, F			
<i>Pieris japonica</i>	Lily-of-the-valley	C, F			
	Mountain fire lily-of-the-valley	C, F			
	Snowdrift lily-of-the-valley	C, F			
	Temple bells lily-of-the-valley	C, F			
	Valley rose lily-of-the-valley	C, F			
	Valley valentine lily-of-the-valley	C, F			
<i>Pieris japonica x forestii</i>	Forest flame lily-of-the-valley	C, F			

<i>Rhododendron</i> <i>hybrid</i> spp.	America rhododendron	C, F	<i>Solanum rantonetii</i>	Paraguay nightshade	C, F
	English Roseum rhododendron	F	<i>Spiraea bumalda</i>	Anthony Waterer spiraea	C, F
	Nova Zembla rhododendron	C, F	<i>Spiraea x cinerea</i>	First snow spiraea	
	Scintillation rhododendron	C, F	'Grefsheim'		
<i>Rhododendron</i> <i>impeditum</i>	Rhododendron	C, F	<i>Spiraea japonica</i>	Dolchia spiraea	C, F
				Japanese alpine spiraea	C, F
<i>Rhododendron indica</i>	Formosa azalea	C, F		Shirobana spiraea	C, F
<i>Rhododendron indica</i>	Waucubusa azalea	C, F	<i>Spiraea vanhouttei</i>	Bridal wreath	C, F
<i>Rhododendron</i> <i>kaempferi</i>	Blue danube azalea	C, F	<i>Syringa rothomagensis</i>	Chinese lilac	C, F
			<i>Syringa vulgaris</i>	Lilac, common	F
<i>Rhododendron kerume</i>	Coral bells azalea	C, F	<i>Taxus cuspidata</i>	Yew, Japanese	F
	Hino crimson azalea	C, F	<i>Tecomaria capensis</i>	Cape honeysuckle	C, F
	Hino pink azalea	C, F	<i>Ternstroemia</i>	Ternstroemia, Japanese	C, F
	Mildred azalea	C, F	<i>gymnanthera</i>		
	Snow azalea	C, F	<i>Thuja occidentalis</i>	Emerald arborvitae	C, F
<i>Rhododendron</i> <i>maximum</i>	Rhodie max (rosebay)	C, F		Globosa-globe arborvitae	C, F
				Little giant-dwarf arborvitae	C, F
<i>Rhododendron</i> <i>mucronulatum</i>	Rhododendron	F		Nigra-dark American arborvitae	C, F
				Pyramidalis arborvitae	C, F
<i>Rhododendron obtusum</i>	Hino-crimson azalea	C, F		Rheingold arborvitae	C, F
<i>Rhododendron</i> <i>ponticum</i>	Chioniodes rhododendron	C, F	<i>Thuja occidentalis</i>	Techny arborvitae	F
			<i>Thuja orientalis</i>	Woodwardii arborvitae	C, F
<i>Rhododendron</i> <i>ponticum</i>	Daphnoides rhododendron	C, F		Aureus nana-dwarf golden arborvitae	C, F
				Minima glauca-dwarf arborvitae	C, F
<i>Rhododendron x</i> <i>'purple gem'</i>	Purple gem rhododendron	C, F	<i>Veitchia merrilli</i>	Christmas palm	F
			<i>Viburnum bodnantense</i>	Pink dawn viburnum	C, F
<i>Rhododendron</i> <i>racemosum</i>	Dwarf scarlet wonder rhododendron	C, F	<i>Viburnum carlesii</i>	Koreanspice viburnum	C, F
	Tribly rhododendron	C, F	<i>Viburnum davidii</i>	David viburnum	C, F
	Unique rhododendron	C, F	<i>Viburnum japonicum</i>	Viburnum	F
	Vulcan rhododendron	C, F	<i>Viburnum judd</i>	Viburnum	C, F
	Ramapo rhododendron	C, F	(v. <i>X juddii</i>)		
<i>Rhododendron</i> <i>sassthigiatim x</i> <i>carolinianum</i>			<i>Viburnum lantana</i>	Wayfaring tree	F
			<i>Viburnum opulus sterile</i>	Common snowball viburnum	F
<i>Rhododendron satuski</i>	Gumpo pink azalea	C, F	<i>Viburnum plicatum</i>	Doublefile viburnum	C, F
	Higasa azalea	F	<i>tomentosum</i>		
		C, F	<i>Viburnum setigerum</i>	Tea viburnum	F
	Reijn azalea	C, F	<i>Viburnum tinus</i>	Spring bouquet viburnum	F
<i>Rhododendron</i> spp. hybrids	Carror azalea	C, F	<i>compactum</i>		
	Fashion azalea		<i>Viburnum trilobum</i>	Cranberry bush	C, F
	Gerard Christina azalea	F	<i>Viburnum trilobum</i>	Dwarf cranberry bush	C, F
	Girard Roberta azalea	C, F	<i>compactum</i>		
	Golden flare exbury azalea	F	<i>Viburnum x pragense</i>	Viburnum	C, F
	Helmut vogel azalea	F	<i>Weigela florida</i>	Bristol ruby weigela	C, F
	Hersey red azalea	F		Java red weigela	C, F
	Hot shot azalea	C, F		Minuet weigela	C, F
	Hume azalea	F	<i>Xylosma congestum</i>	Xylosma	F
	Inga azalea	F	<i>Yucca filamentosa</i>	Yucca	C, F
	Irene Koster azalea	C, F			
	President Clay azalea	C, F			
	Tradition azalea	C, F			
<i>Rhus lancea</i>	Sumac, African	C, F			
<i>Rosa rugosa</i>	Ramanas rose	C, F			
<i>Rosmarinus officinalis</i>	Rosemary	F			
<i>Senecio cineraria</i>	Dusty-miller/silver ragweed	C, F			
<i>Skimmia japonica</i>	Japanese skimmia	C, F			
<i>Skimmia revesiana</i>	Reeve's skimmia	C, F			

**Groundcovers/
Perennials**

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name				
<i>Achillea millefolium</i>	Common yarrow	C, F	<i>Fuchsia x 'Santa Claus'</i>	Santa Claus Fuchsia	C, F
<i>Agapanthus africanus</i>	Lily of the Nile	C, F	<i>Gaillardia aristata</i>	Blanket flower	C, F
<i>Agapanthus "Peter Pan"</i>		C, F	<i>Gaillardia grandiflora</i>	Goblin blanket flower	C, F
<i>Alstroemeria aurea</i>	Peruvian lily	C, F	<i>Gaura lindheimeri</i>	Gaura	C, F
<i>Ammophila breviligulata</i>	Beechgrass	C, F	<i>Gazania rigens</i>	Gazania, trailing	C, F
<i>Antirrhinum majus</i>	Snapdragon	C, F	<i>leucolaena</i>		
<i>Arctotheca calendula</i>	Cape weed	F	<i>Gazania spp.</i>	Gazania	C, F
<i>Argyranthemum frutescens</i>	Paris daisy	C, F	<i>Geranium incanum</i>	Cranesbill	C, F
<i>Artemisia schmidtiana</i>	Angels' hair	C, F	<i>Geranium subcaulescens</i>	Black eyed magenta cranesbill	C, F
<i>Asparagus retrofractus</i>	Fern	C, F	<i>Hakonechloa macroaureola</i>	Golden hakonechloa	C, F
<i>Asteriscus maritimus</i>	Gold coin daisy	C, F	<i>Hedera canariensis</i>	Ivy, Algerian	F
<i>Astilbe Deutschland</i>	Deutschland astilbe	C, F	<i>Hedera helix</i>	Ivy, English	C, F
<i>Asparagus retrofractus</i>	(No common name)	C, F	<i>Helichrysum petiolatum</i>	White licorice plant	C, F
<i>Asparagus variegata</i>	Tree fern	C, F	<i>Hemerocallis spp.</i>	Daylily	C, F
<i>Aster novae-angliae</i>	New England aster	C, F	<i>Hesperaloe parviflora</i>	Red yucca	C, F
<i>Aster novi-belgii</i>	New York aster	C, F	<i>Heuchera americana</i>	Palace purple	C, F
<i>Begonia cordifolia</i>	Heartleaf begonia	C, F	<i>Heuchera micrantha</i>	Coral bells	C, F
<i>Begonia semperflorens</i>	White ambassador begonia	C, F	<i>Hippeastrum hybrid</i>	Amaryllis	C, F
<i>Bidens ferulifolia</i>	Peter's Gold Bidens	C, F	<i>Hosta 'Francee'</i>	Francee plantain lily	C, F
<i>Brachycome x 'New amethyst'</i>	Swan River Daisy New Amethyst	C, F	<i>Hosta lancifolia</i>	Albo-marginata hosta	C, F
<i>Callistephus chinensis</i>	China aster	C, F	<i>Hosta 'Patriot'</i>	Patriot plantain lily	C, F
<i>Carex spp.</i>	Variiegated carex	C, F	<i>Hymenoxys acaulis</i>	Angelita daisy	C, F
<i>Carpobrotus edulis</i>	Ice plant, largeleaf (see label)	F	<i>Hypericum spp.</i>	St. Johnswort	C, F
<i>Catharanthus roseus</i>	Madagascar periwinkle	C, F	<i>Impatiens wallerana</i>	Busy lizzie	C, F
<i>Cerastium tomentosum</i>	Snow in the summer	C, F	<i>Iris pumila</i>	Yellow dwarf bearded iris	C, F
<i>Ceratostigma plumbaginoides</i>	Dwarf plumbago	C, F	<i>Iris siberica</i>	Blue siberian iris	C, F
<i>Chrysanthemum morifolium</i>	Florist's chrysanthemum	C, F	<i>Jasminum nitidum</i>	Angelwing jasmine	C, F
<i>Chrysanthemum sp.</i>	Chrysanthemum species	C, F	<i>Lampranthus spectabilis</i>	Trailing iceplant	F
<i>Clematis integrifolia caerulea</i>	Blue bell clematis	C, F	<i>Leptospermum scoparium</i>	Broom teatree/manuka	C, F
<i>Clivia miniata</i>	Kafir lily	C, F	<i>Liatris spicata</i>	Gay feather	C, F
<i>Coreopsis verticillata</i>	Coreopsis, threadleaf	C, F	<i>Limonium perezii</i>	Statice	C, F
<i>Cortaderia selloana</i>	Pampas grass	C, F	<i>Liriope gigantea</i>	White lily turf	C, F
<i>Cuphea hyssopifolia</i>	False or Mexican heather	C, F	<i>Liriope muscari</i>	Lilac beauty lily turf	C, F
<i>Cyperus albostratus</i>	Dwarf umbrella grass	C, F		Majestic lily turf	C, F
<i>Dahlia x 'Royal dahlia pink'</i>	Dwarf dahlia Wendy pink	C, F	<i>Liriope spicata</i>	green/creeping lily turf	C, F
<i>Delosperma alba</i>	White iceplant	F	<i>Lobelia erinus</i>	Lobelia	C, F
<i>Descampsia caespitosa</i>	Descampsia	C, F	<i>Lobularia maritima</i>	Sweet alyssum	C, F
<i>Dianthus gratianopolitanus</i>	Crimson treasure cheddar pink	C, F	<i>Lonicera japonica</i>	Honeysuckle, Japanese	F
<i>Dietes vegeta</i>	Fortnight lily	C, F	<i>Lysimachia punctata</i>	Dotted loosestrife	C, F
<i>Drosanthemum floribundum</i>	Trailing rosea iceplant	F	<i>Mathiola incana</i>	Stock	C, F
<i>Drosantheumum hispidum</i>	Iceplant	C, F	<i>Miscanthus sinensis</i>	Eulalia grass	C, F
<i>Ensete ventricosum</i>	Red abyssinian banana	C, F	<i>Monarda didyma</i>	Bee balm	C, F
<i>Equisetum scirpoides</i>	Dwarf horsetail	C, F	<i>Moraea iridiodes</i>	African iris	C, F
<i>Erianthus ravennae</i>	Hardy pampasgrass	C, F	<i>Oenothera speciosa</i>	Siskiyou evening primrose	C, F
<i>Erysimum "Bowles mauve"</i>	Wallflower	C, F	<i>Ophiopogon japonicus</i>	Dwarf Mondo grass	C, F
<i>Euryops pectinatus</i>	Dwarf euryops	C, F		Mondo grass	C, F
<i>Eustoma grandiflorum</i>	Pink lisianthus	C, F	<i>Osteospermum fruticosum</i>	Freeway daisy	C, F
<i>Festuca ovina glauca</i>	Blue fescue	C, F	<i>Pachysandra terminalis</i>	Japanese spurge	C, F
			<i>Parthenocissus quinguefolia</i>	Virginia creeper	C, F
			<i>Pelargonium x hortorum</i>	Zonal geranium	C, F
			<i>Pelargonium peltatum</i>	Ivy geranium	C, F
			<i>Pennisetum alopecuroides</i>	Fountain grass	C, F

<i>Pennisetum setaceum</i>	Chrimson fountaingrass	C, F
<i>Pentas lanceolata</i>	Star cluster	C, F
<i>Penstemon x</i> 'Apple blossom'	Apple blossom penstemon	C, F
<i>Penstemon</i> <i>gentianoides</i>	Hartwig penstemon	C, F
<i>Perovskia atriplicifolia</i>	Russian sage	C, F
<i>Petunia-hybrids</i>	Garden petunias	C, F
<i>Phalaris arundinacea</i> <i>picta</i>	Ribbon grass	C, F
<i>Ratibida columnifera</i>	Mexican hat	C, F
<i>Rudbeckia fulgida</i>	Blackeyed Susan	C, F
<i>Rudbeckia hirta</i>	Blackeyed Susan	C, F
<i>Ruellia brittoniana</i>	Dwarf Katie ruellia	C, F
<i>Salvia grahamii</i>	Graham's sage	C, F
<i>Salvia leucantha</i>	Mexican bush sage	C, F
<i>Sedum x 'Autumn joy'</i>	Autumn joy stonecrop	C, F
<i>Sedum x 'Vera jameson'</i>	Vera jameson stonecrop	C, F
<i>Targetes patula</i> 'Little hero'	Little hero marigold	C, F
<i>Trachelospermum</i> <i>asiaticum</i>	Asian jasmine	C, F
<i>Tulbaghia violacea</i>	Society garlic	C, F
<i>Verbena peruviana</i>	St. Paul verbena	C, F
<i>Vinca major</i>	Periwinkle, bigleaf	C, F
<i>Vinca minor</i>	Periwinkle, dwarf	F
<i>Vinca</i> spp.	Periwinkle	F
<i>Zinnia elegans</i>	Dwarf zinnia	C, F

pecan	F
pistachio	F
plum	F
pomegranate	F
prune	F
raspberry	F
walnut, black	F
walnut, English	F
†Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.	

Ornamental Bulbs

Do not use Snapshot 2.5 TG in bulb production fields. For use in landscape settings only.

Snapshot 2.5 TG may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply Snapshot 2.5 TG to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. Snapshot 2.5 TG may also be applied following bulb emergence but prior to flowering and after flowering. For fall planted bulbs, apply Snapshot 2.5 TG again in late winter or early spring to weed-free soil surfaces.

Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.

Do not apply to gladiolus prior to emergence or if corms are less than one inch in diameter.

Do not apply to bulbs while they are flowering.

Shadehouse Areas

Snapshot 2.5 TG may be applied in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within three weeks prior to enclosing greenhouse or poly-type structures.

Noncropland

Snapshot 2.5 TG is recommended as a preemergence treatment for control of certain broadleaf weeds and annual grasses on non-cropland areas such as industrial sites, utility substations, highway guardrails, sign posts, delineators, etc. Apply Snapshot 2.5 TG anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the "General Information" section of this label for General Use Precautions and information on Application, Application Rates, and Weeds Controlled.

Non-bearing Fruit and Nut Tree and Non-bearing Vineyards†

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Common Name

almond	F
apple	F
apricot	F
avocado	F
blackberry	F
blueberry	F
boysenberry	F
cherry, sour	F
cherry, sweet	F
currant	F
dewberry	F
elderberry	F
fig	F
filbert	F
gooseberry	F
grape, American	F
grape, European	F
grapefruit	F
kiwi	F
lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive	F
orange	F
peach	F
pear	F

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Label Code: D02-082-016

Replaces Label: D02-082-015

EPA-Accepted 03/20/2002

Revisions:

Label Changes by Amendment EPA-accepted June 19, 2000:

1. **Precautionary Statements:** Revised First Aid statement text in accordance with new EPA requirements.
2. **Directions for Use:** (1) Added comment regarding accidental application to turfgrasses adjacent to ornamental plantings. (2) Expanded listing of ornamental plant species (39 new tree, shrub and perennial/ground cover species added) in which Snapshot 2.5 TG may be applied at labeled rates.